

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/305446021>

Characterization of best naringinase producing fungus strain isolated from palmyrah (*Borrasus flabellifer*) fruit pulp

Article in *International Journal of Biological Research* · July 2016

DOI: 10.14419/ijbr.v4i2.6289

CITATIONS

4

READS

319

1 author:



Ranganathan Kapilan
University of Jaffna

81 PUBLICATIONS 383 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Biodiesel production from marine flora and optimized the condition for yield enhancement [View project](#)



microbial protein [View project](#)



Characterization of best naringinase producing fungus strain isolated from palmyrah (*Borrassus flabellifer*) fruit pulp

Sinthuja Karuppaija^{1*}, Ranganathan Kapilan², Vasantharuba Seevaratnam¹

¹Department of Agricultural Chemistry, Faculty of Agriculture, University of Jaffna, Sri Lanka

²Department of Botany, Faculty of Science, University of Jaffna, Sri Lanka

*Corresponding author E-mail: ranganat@ualberta.ca

Abstract

Background: The Palmyrah (*Borrassus flabellifer* L.) fruit pulp has the bitter compound flabelliferin (a tetraglycoside) which can be hydrolyzed by naringinase enzyme. The diverse groups of filamentous fungi and bacteria that live in different substrates have the capacity of producing extracellular naringinase enzyme which is of tremendous industrial value.

Objective: The objective of the study was to isolate the naringinase producing fungal strains from Palmyrah and to identify the best naringinase producer under liquid and solid state fermentation systems.

Methods: Fungal strains isolated from Palmyrah fruit pulp and the soil where pulp is allowed to decay, were grown on naringin agar selective medium at pH 6.0 at room temperature and the production of extracellular naringinase was measured in the liquid fermentation media and solid state fermentation system using paddy husk as support.

Results: Five fungal strains isolated from the palmyrah pulp and the pulp decaying in sand designated as PF1, PF2, PF3, PF4 & PF5 had the ability to produce extracellular naringinase enzyme in liquid fermentation media. Fungal strain PF4 that showed highest naringinase enzyme activity (1.769U/ml) was selected among the isolated five fungal strains and identified as *Rhizopus stolonifer* based on the morphological and biochemical characteristics. When this strain was grown in the solid state fermentation system using paddy husk as media, naringinase production was higher (269.84 U/gram of dry substrate) in seven days.

Conclusion: *Rhizopus stolonifer* could be used to produce large scale naringinase enzyme under solid state fermentation system using very cheap, easily available, agricultural waste paddy husk as support without the need of expensive and well equipped laboratories.

Keywords: Naringinase; Paddy Husk; Palmyrah Pulp; *Rhizopus stolonifer*.

1. Introduction

Palmyrah (*Borrassus flabellifer* L.) is one of the abundant palm in the dry zone of the northern and southern part of Sri Lanka. It is said as “the life of tree” because almost all part of the tree has economic value. Potentially a lot of food and beverage items are prepared from the Palmyrah fruit pulp for the commercial purpose in the developing countries like, India, Pakistan, Bangladesh and Sri Lanka. Even though the demand of the food products is substantially high, a certain percentage of consumers complain about the bitterness of the fruit pulp that lead to the unacceptable quality. Due to this reason, more than half of the annual production of Palmyrah fruits are thrown away and used as animal feed (Jansz et al, 2002). Traditionally a lot of methods like heat treatments and chemical methods are in practice to reduce the bitterness of the food and beverages of the Palmyrah pulp but these methods are less effective and economically unfavorable.

The bitter compound that present in the palmyrah fruit pulp is flabelliferin (Fii) which is identified as a tetraglycoside (Nikawela et al, 2000, Nikawela et al, 2011 and Jayaratnam, 2015). The flabelliferin (Fii) can be hydrolyzed into rhamnose and glucose molecules and 40% of bitterness of flabelliferin (Fii) can be hydrolyzed by α -amylase and some are not susceptible to α -amylase (Jansz et al, 2002 and Zhang et al, 2007). All the bitter compound of flabelliferin (Fii) can be hydrolyzed by the glycolytic enzyme naringinase (Jansz et al, 2002 and Jayaratnam 2015). The nar-

inginase is a complex enzyme which has α -L-rhamnosidase and β -D-glucosidase activities that leads to potential usefulness in pharmaceuticals and food industries (Ribeiro 2011). In recent years, naringinase enzyme is widely used in debittering of palmyrah cordial, jams and citrus juice industries (Jansz et al, 2002 and Kumar et al, 2010). The naringinase enzyme can be produced naturally from plants, fungus and bacteria. There have been some studies on the filamentous fungi like *Penicillium* sp, *Aspergillus* sp, *Rhizopus* sp and those studies confirmed that these organisms have the ability to produce naringinase enzyme (Mendoza-Cal 2010) when the growth substrate contains bitter compound like Naringin (flabelliferin).

In the case of naringinase production at the commercial level, the filamentous fungus might take an ideal option for their cost effective and economically viable in nature than the chemical methods of naringinase production. Since there have been no reports about isolating a naringinase producer from underutilized Palmyrah pulp and its decay in soil, it was planned to work on this area. The objective of the study was to isolate naringinase producing fungal strains from the fruit pulp of palmyrah and its decay in soil and to characterize the best naringinase producer under liquid and solid fermentation system.

2. Materials and methods

2.1. Source of organisms



Decaying palmyrah fruit pulp and buried fruit pulp in the soil for two weeks, were the sources of organisms.

2.2. Chemicals

Naringin, Potato dextrose agar(PDA), Glucose, Peptone, Yeast extract, KH_2PO_4 , $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, DNS solution

2.3. Isolation of naringinase producing fungus

The isolation of naringinase producing fungus was carried out as follows: The decaying fruit pulp and soil buried fruit pulp were taken to make serial dilution up to 10^{-6} and 1ml was inoculated on 0.1% (w/v) naringin and 0.5% (w/v) glucose agar medium but the soil sample around buried fruit was taken and mixed with sterile distilled water and the filtrate was taken to dilute until 10^{-4} from diluted sample, 1ml was inoculated on the above medium. After 3 days, based on the morphological observations, every fungal strains were sub-cultured on the freshly reared medium until pure cultures were obtained. Five different fungal strains with different morphological and cultural features were obtained. These five fungal strains (labeled as PF1, PF2, PF3, PF4 & PF5) were inoculated on 0.2% (w/v) naringin and 0.2% (w/v) glucose agar medium to screen naringinase producing fungus based on the colour change of the medium. Secondary screening was also conducted by testing the naringinase enzyme by the assay method (Sinthuja et al. 2016). The isolated fungal strains were sub-cultured on PDA slant and stored at 4°C .

2.4. Fermentation

2.4.1. Liquid state fermentation

The fermentation medium containing (w/v) 0.2% naringin, 0.5% glucose, 0.7% peptone, 0.1% yeast extract, 0.05% KH_2PO_4 , 0.01% $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ and 0.07% $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 0.07% $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.07% $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ was used. The pH was adjusted to 6.0 (Navaratnam et al. 2003). Twenty five mL of the sterilized liquid medium was taken into 100ml conical flask and 6day old cultures of each organism from potato dextrose agar (PDA) medium was inoculated in the liquid fermentation medium and allowed for fermentation at 200rpm in room temperature until 9th day. The spores were taken out using sterile 0.85% NaCl solution and spore size was adjusted as 10^8 spores/ml. 1ml of spore solution was inoculated on duplicate fermentation medium of each fungal strain under aseptic condition. The fermented sample was withdrawn for the enzyme assay at 7th, 8th and 9th day. One unit of naringinase activity is defined as the amount of enzyme that produces $1\mu\text{mol}$ of reducing sugar in 1 minute at pH 5 and 60°C with 10gL^{-1} naringin.

$$\text{Enzyme Activity} = \frac{\text{concentration of glucose standard(mg)} \times \text{absorbance of test} \times 1000}{\text{Absorbance of standard glucose} \times \text{volume of enzyme solution(ml)} \times \text{incubation time(min)} \times 180}$$

2.5.2. Solid fermented sample

The culture filtrate was taken from solid state fermentation medium after 7th day to find out enzyme activity. Enzyme assay was done by modified Miller method as same as liquid fermented sample and 1gram of solid substrate was taken from solid fermentation medium to analyze the moisture content which was carried out at 80°C until getting a constant weight. The enzyme activity of solid state fermentation was presented as unit (U) per gram of dry substrate.

2.4.2. Solid state fermentation

The solid state fermentation was carried out for selected to compare the enzyme activity with liquid fermentation because of limited facilities and use of less expensive agricultural waste materials to make cost effective production. Fermentation medium contained 2:10 ratio of paddy husk and liquid medium. Twenty grams of paddy husk was inoculated with 1ml of the spore suspension with 10^8 spores/ml of the selected fungal strain *Rhizopus stolonifer* in 100ml liquid medium at pH 6.0 and allowed to ferment for 7 days in room temperature. Sample was withdrawn and filtered and the filtrate was used for the enzyme assay. Duplicate was also maintained.

2.5. Enzyme assay

Enzyme assay was done based on the modified Miller (1959) method developed by Roy and Uddin (2004). Naringinase enzyme has α -L-rhamnosidase and β -D-glucosidase that hydrolysis flabeliferin into glucose as end product. The glucose is a reducing sugar that has aldehyde group. This group reduces 3,5-dinitrosalicylic acid (DNS) to orange coloured 3-amino, 5 nitrosalicylic acid in alkaline medium on boiling and oxidizes itself to carboxylic group. Sodium potassium tartarate helps to fix the colour of 3-amino 5-nitrosalicylic acid and intensity of the orange colour developed is proportional to the concentration of reducing sugar present in the solution. Based on this principle, the activity of naringinase was determined by glucose standard curve.

2.5.1. Liquid fermented sample

The sample was withdrawn from liquid fermentation medium after 7th, 8th and 9th day of incubation. The sample was centrifuged at 10000rpm for 20 minutes. The supernatant was taken as crude enzyme for determination of naringinase activity. Duplicates and controls were also kept for each sample. 0.25ml of substrate that contain 0.5% naringin at pH 5.0 in citrate buffer was taken and allowed for pre-incubation for 3min at 50°C and simultaneously enzyme also was subjected to pre incubation. Then 0.25ml of enzyme (supernatant) was added to each test except control. It was allowed to reaction at 50°C for 10min. Then 0.5ml of DNS solution was added immediately to stop the reaction. Finally the container was boiled and cooled and then 5ml of sterilized distilled water was added and the volume was made to 6ml. After 3min of pre incubation 0.5ml of DNS solution was added then boiled and it was allowed to cool and after that 5ml of sterilized distilled water was added and this set up was used as control. Test and controls were stirred before the reading by using spectrophotometer at 550nm (Navaratnam et al, 2003, Sinthuja et al. 2016, Keerthini et al. 2016).

Unit of enzyme activity (U/ml): Amount of naringin (μ mole) which can be hydrolyzed by 1ml of crude enzyme naringinase in 1 minute.

Enzyme activity was determined by the following equation,

2.6. Identification of fungi

Potato dextrose agar (PDA) slant and plates samples were prepared and maintained at 25°C for identification of selected fungal strain. Morphological characteristics were used to identify the fungi. The colony colour, sporangial shape and structures were observed from initial stage to maturity with naked eye and through microscope. The sporangiospore, Rhizoids, stolons and sporan-

giophore were also observed under microscope. The spore size was measured with the help of stage microscope.

3. Results and discussion

3.1. Isolation of naringinase producing strains

In this study, natural Palmyrah wastes such as palmyrah pulp and decaying particles of the pulp in the sand were selected as source for the isolation of naringinase producing fungus. Primary screening was done by observation of culture medium. Five fungal strains / colonies changed the colour of the selective medium.




Fig. 1: Isolated Fungal Strains.

Table 1: Enzyme Activity of Isolated Fungal Strains in Liquid Fermentation Method

Organism	6 th Day Activity (U/ml)	7 th Day Activity (U/ml)	8 th Day Activity (U/ml)
PF1	1.536	1.653	1.375
PF2	1.327	1.075	0.877
PF3	1.701	1.711	0.931
PF4	1.769	1.747	1.359
PF5	1.023	1.055	0.6

Naringin agar selective medium was used for the primary screening. The best naringinase producing strain was isolated from palmyrah fruit fiber and 1.656U/ml of activity was obtained at pH 5.0 in citrate buffer and at 50°C (Navaratnam et al, 2003). The activity of the selected strain was higher than previous organism isolated. The difference in the amount of enzyme production may be due to the variation of fungal strains, source and cultural conditions.

The easily available Palmyrah fruit pulp was selected as source to isolate the naringinase producing fungal strains and the natural diversity is important to adopt the organism to produce appropriate enzyme (Thammawat et al, 2008). In this study decayed palmyrah fruit pulp, fruit pulp that is buried in the soil and soil sample around the buried pulp were used to isolate the fungal strains (Table 2). Palmyrah fruit fiber was selected to isolate naringinase producing strains and 25 fungal strains were isolated from palmyrah fruit fiber as naringinase producing strains (Navaratnam 2003). *Aspergillus flavus* isolated from the wet soil in a poultry house produced naringinase (Radhakrishnan and Sampath, 2013). *Aspergillus niger* BCC 25166 was proved to produce naringinase enzyme (Thammawat et al, 2008).

Table 2: Isolated Fungal Strains in this Study

Fungal Strains	Source
PF1	Soil Around Buried Fruit Pulp
PF2	Buried Fruit Pulp
PF3	Decayed Fruit Pulp
PF4	Decayed Fruit Pulp
PF5	Buried Fruit Pulp

Penicillium decumbens PTCC 5248 was reported as naringinase producing strain in liquid media when cultivated in medium containing naringin as carbon source (Norouzian et al, 2000). The carbon source could be considered as inducer for naringinase pro-

duction. Secondary screening was done by measuring the naringinase enzyme production. It was five fungal strains from primary screening were selected as naringinase producers. Those strains were capable of producing extracellular naringinase enzyme in liquid fermentation media under no agitation. Among the isolated fungal strains, only five fungal strains were identified as naringinase producers and designated as PF1, PF2, PF3, PF4 & PF5 - Fig 1). Higher enzyme activity was observed on the 7th day for PF1 and PF4 and 8th day for PF2, PF3 and PF5 (Table 1). PF4 showed highest naringinase activity (1.769U/ml) at pH 5.0 in citrate buffer when the temperature was kept at 50°C.

The naringin was the best inducer for the production of naringinase by *Aspergillus niger* MTCC 1344 than hesperidin, naringenin, rhamnase and rutin. Enzyme activity was optimum at 7th day of incubation (Kumar 2015). Most *Aspergillus* strains (*A. foetidus*, *A. niger* HPD2, *A. niger*) showed higher enzyme activity on the 8th day of incubation (Mendoza-Cal, 2010). Naringinase enzyme activity was higher on the 7th day of incubation.

3.2. Solid state fermentation of selected strain

When the selected fungal strain (PF4) was subjected to solid state fermentation using paddy husk as support, highest naringinase enzyme activity (269.84 U/gram of dry substrate) was obtained on the 7th day at room temperature under no agitation.

There have been no reports available for the production of naringinase by *Rhizopus stolonifer* by using paddy husk as substrate in solid state fermentation but potential to have produce enzyme naringinase by filamentous fungi in solid state fermentation. The solid state fermentation process was best method over conventional submerged fermentation because of higher yield of secondary metabolites and enzymes. Certain enzymes and secondary metabolites can only be produced under solid state fermentation system (Barrios-González and Tarragó-Castellanos, 2015). The production of any compound by using microorganism must be considered with the facilities, amount of outcome, cost effectiveness and also satisfaction of the compound characteristics (Pandey et al, 1996). The growth and metabolic activity of the microorganism can be facilitated by solid substrate which absorbs the moisture and provides better utilization of substrate by microorganisms (Pandey 1992).

The naringinase enzyme was produced by filamentous fungus using grape fruit rind and orange rind as substrates under solid

state fermentation and the grape fruit rind was best substrate than orange rind (Mendoza-Cal 2010). The *Aspergillus niger* MTCC 1344 produced high levels of naringinase enzyme in a short period about 96 hours of incubation when solid state fermentation with rice bran was done (Ni et al, 2011). The production of α -L-rhamnosidase by *Aspergillus terreus* CECT 2663 was highest in solid state fermentation using sugar cane bagasse as substrate (Elinbaum et al, 2002). Similarly different naringinase producing fungal types from different species were isolated from citrus maxima fruits (Keerthini et al, 2016 a, b) and decaying Palmyrah fruit pulp in soil (Sinthuja et al, 2016). Glucoamylase production by *Aspergillus niger* CFTRI 1105 was highest when using paddy husk as support (Arasaratnam et al, 2001). Xylanase production by *Bacillus pumilus* in solid state fermentation using paddy husk as support yielded significantly higher production of enzyme than the usual liquid media (Kapilan and Arasaratnam, 2011). The *Rhizopus stolonifer* was used to produce cellulose-free alkaline xylanase in solid state using wheat bran as substrate (Goulart 2005). Diverse studies indicate that potential and possibilities to get commercial level of enzyme (naringinase) production by filamentous fungus using solid state fermentation (paddy husk).

3.3. Identification of selected fungal strain




Fig. 2: Sporangiospores of *Rhizopus Stolonifer* (A), Growth of *Rhizopus Stolonifer* under Microscope (B), Under Naked Eye (C) on Potato-Dextrose Agar (PDA) Medium.

4. Conclusion

Among the five naringinase producing fungal strains isolated from Palmyrah pulp, the strain that was confirmed as *Rhizohpus stolonifer* showed highest naringinase enzyme activity in the liquid fermentation system (1.769U/ml) and in the solid state fermentation system using paddy husk asa support (269.84U/gram of dry substrate). Solid phase production of naringinase with the support of paddy husk for this fungus would be an efficient cost effective method for the production of large scale naringinase.

Acknowledgement

Authors sincerely thank the University of Jaffna for awarding the Research Grant to Dr.S.Vasantharuba, Department of Agricultural Chemistry. Technical support and suggestions provided by Mr. V. C. Anpalagan, Mr. P.T.J. Jashothan and Mr.A.Thananchayan, University of Jaffna is highly appreciated.

References

- [1] Arasaratnam, V., Mylvaganam, K., &Balasubramaniam, K. (2001). Glucoamylase production by *Aspergillusniger* in solid state fermentation with paddy husk as support. *Journal of food science and technology*, 38(4), 334-338.
- [2] Barrios-González, J., and Tarragó-Castellanos, M. R. (2015). Solid-State Fermentation: Special Physiology of Fungi.
- [3] Bram, B., &Solomons, G. L. (1965). Production of the enzyme naringinase by *Aspergillusniger*. *Applied microbiology*, 13(6), 842-845.
- [4] CO, O. (2011). *Rhizopusstolonifer* exhibits dimorphism. *Afri. J. Biotechnol*, 10 (20), 4269-4275.
- [5] Elinbaum, S., Ferreyra, H., Ellenrieder, G., & Cuevas, C. (2002). Production of Aspergillusterreus β -L-rhamnosidase by solid state fermentation. *Letters in applied microbiology*, 34(1), 67-71. <http://dx.doi.org/10.1046/j.1472-765x.2002.01039.x>.
- [6] Goulart, A. J., Carmona, E. C., &Monti, R. (2005). Partial purification and properties of cellulase-free alkaline xylanase produced by *Rhizopusstolonifer* in solid-state fermentation. *Brazilian Archives of Biology and Technology*, 48 (3), 327-333. <http://dx.doi.org/10.1590/s1516-89132005000300001>.
- [7] Hernández-Lauzardo, A. N., Bautista-Baños, S., Velázquez-del Valle, M. G., & Trejo-Espino, J. L. (2006). Identification of *Rhizopusstolonifer* (Ehrenb: Fr.) Vuill, causal agent of Rhizopus rot disease of fruits and vegetables. *Revista Mexicana de Fitopatología*, 24(1), 65-69.
- [8] Jansz, E. R., Nikawela, J. K., Gooneratne, J., &Thevendirarajah, K. (1994). Studies on the bitter principle and debittering of Palmyrah fruit pulp. *Journal of the Science of Food and Agriculture*, 65(2), 185-189. <http://dx.doi.org/10.1002/jsfa.2740650210>.
- [9] Jayaratnam, M. (2015). *The chemistry and biochemistry of palmyrah products* (Doctoral dissertation).
- [10] Kapilan, R., and Arasaratnam, V. (2011). Paddy husk as support for solid state fermentation to produce xylanase from *Bacillus pumilus*. *Rice Science*, 18 (1), 36-45. [http://dx.doi.org/10.1016/S1672-6308\(11\)60006-1](http://dx.doi.org/10.1016/S1672-6308(11)60006-1).
- [11] Keerthini, S., Kapilan, R., Vasantharuba, S. (2016a). Characterization of best naringinase producing fungus isolated from the citrus fruits. *International Journal of Biological Research*, 4(2), 83-87. <http://dx.doi.org/10.14419/ijbr.v4i2.6281>.

- [12] Keerthini, S, Kapilan, R., Vasantharuba, S. (2016b). Optimization of naringinase production from *Aspergillus flavus* in solid state fermentation media using citrus peel as support, *Scholars Academic Journal of Biosciences*, 4 (6), 535-543. DOI: 10.21276/sajb.2016.4.6.16
- [13] Kumar, V. V. (2015). Comparative studies on inducers in the production of naringinase from *Aspergillus niger* MTCC 1344. *African Journal of Biotechnology*, 9(45), 7683-7686.
- [14] Kwon, J. H., Kang, S. W., Kim, J. S., & Park, C. S. (2001). Rhizopus soft rot on cherry tomato caused by *Rhizopus stolonifer* in Korea. *Mycobiology*, 29(3), 176-178.
- [15] Mendoza-Cal, A., Cuevas-Glory, L., Lizama-Uc, G., & Ortiz-Vázquez, E. (2010). Naringinase production from filamentous fungi using grapefruit rind in solid state fermentation. *African Journal of Microbiology Research*, 4(19), 1964-1969.
- [16] Navaratnam, P., Ketheeswary, N., Senthuran, A., & Arasaratnam, V. (2003). Preliminary studies on the isolation of naringinase producing fungus.
- [17] Ni, H., Li, L., Xiao, A., Cao, Y., Chen, Y., & Cai, H. (2011). Identification and characterization of a new naringinase-producing strain, Williopsis californica Jmudeb007. *World Journal of Microbiology and Biotechnology*, 27(12), 2857-2862. <http://dx.doi.org/10.1007/s11274-011-0766-7>.
- [18] Nikawela, J. K., Abeyssekera, A. M., & Jansz, E. R. (2011). Flabelliferins-steroidal saponins from palmyrah (*Borassus flabellifer* L.) fruit pulp isolation by flash chromatography, quantification and saponin related activity. *Journal of the National Science Foundation of Sri Lanka*, 26(1). <http://dx.doi.org/10.4038/jnsfstr.v26i1.3080>.
- [19] Nikawala, J. K., Jansz, E. R., Baeckstrom, P., Abeyssekera, A. M., & Wijeyaratne, S. C. (2000). Flabelliferins of naringinase debittered palmyrah fruit pulp.
- [20] Norouzian, D., Hosseinzadeh, A., Inanlou, D. N., & Moazami, N. (2000). Production and partial purification of naringinase by *Penicillium decumbens* PTCC 5248. *World Journal of Microbiology and Biotechnology*, 16(5), 471-473. <http://dx.doi.org/10.1023/A:1008962131271>.
- [21] Pandey, A. (1992). Recent process developments in solid-state fermentation. *Process Biochemistry*, 27(2), 109-117. [http://dx.doi.org/10.1016/0032-9592\(92\)80017-W](http://dx.doi.org/10.1016/0032-9592(92)80017-W).
- [22] Radhakrishnan, I., & Sampath, S. Isolation and characterization of enzyme naringinase from *Aspergillus flavus*. *International Journal of Advanced Biotechnology and Research*, 1(4), 208-212.
- [23] Ribeiro, M. H. (2011). Naringinases: occurrence, characteristics, and applications. *Applied microbiology and biotechnology*, 90(6), 1883-1895. <http://dx.doi.org/10.1007/s00253-011-3176-8>.
- [24] Sinthuja, K., Kapilan, R., Vasantharuba, S. (2016). Optimization of naringinase production by *Rhizopus stolonifer* in solid state fermentation media using paddy husk as support, *Scholars Academic Journal of Biosciences*, 4(6), 507-514. DOI: 10.21276/sajb.2016.4.6.10
- [25] Sumuduni, K. A. V., Jansz, E. R, and Wickremasekara, N. T. (2002). A review of the chemistry and biochemistry of seed shoot flour and fruit pulp of the palmyrah palm (*Borassus flabellifer* L.).
- [26] Thammawat, K., Pongtanya, P., Juntharasri, V., & Wongvithoonyaporn, P. (2008). Isolation, preliminary enzyme characterization and optimization of culture parameters for production of naringinase isolated from *Aspergillus niger* BCC 25166. *Kaestsart J Nat Sci*, 42, 61-72.