COMPARATIVE STANDARDIZATION OF SIDDHA MEDICATED OIL: THALANGAI ENNAI, A POLY HERBAL FORMULATION

S. Vinotha 1*, I. Thabrew2, S. Sri Ranjani 3

^{1,3} Unit of Siddha Medicine, University of Jaffna, Sri Lanka
 vsanmuga07@gmail.com
²Institute of Biochemistry, Molecular Biology and Bio technology, University of
 Colombo, Sri Lanka

Key words: Thalangai ennai, Poly herbal formulation, Medicated Oil, Standardization

Introduction

With the 'retain to nature' call in both, the developed countries as well as developing countries, there is an increase in number of people switching over to the alternative system of medicine and thus it is essential that people get relevant medicines (Kumar & Baxi, 2007). Siddha Medicine is a one of the traditional medicine (Thirunarayanan, 1988) and practiced by Tamil people of Northern and Eastern province in Sri Lanka. An attempt is made to standardize Siddha medicated oil.

Effective management of vātā disorders such as rheumatic pain, inflammation, bone degenerative changes and impaired movement of limbs are the important tasks in the field of treatment. Siddhars described vātā disorders of different origin and external application is most useful for their management. Among them Ennai

(Medicated oil) is one of the medicaments prescribed by them.

Thalangai ennal is one of the Siddha medicated oil mentioned in the Siddha Literature (Ponniahpillai, 2000) and it is prepared at Government Indigenous Medicine Drug Unit in Jaffna and supplied to all Government District Ayurvedic Hospital, Rural Ayurvedic Hospitals and Central Ayurvedic Dispensaries in Jaffna District. marketed formulation of Thalangai ennai (Everest product) is available in Jaffna District. The ingredients used in Thalangai ennai are juice of the thalangai (Pandanus tectorius), sesame oil, milk, water, rock salt and seventeen different herbal powdered materials with that it is one of the most effective oil for external applications.

The physicochemical standards available for the standardization of Siddha medicated oils are insufficient. The analytical values available in the

pharmacopoeial standards for siddha formulations are not finger print standards for each of oil (Hepsibah et al. 1998). The aim of the present study was fixed and compares preliminary reference standards of the Thalangai ennai. This article reports on comparative preliminary standardization of Thalangai ennai (preparation of Drug Unit, Indigenous Medicine and the marketed formulation) have been standardized on the basis of physicochemical properties, organoleptic characters, and Thin Layer Chromatography (TLC) finger print analysis.

Methodology Marketed samples for external medicine

Thalangai ennai was purchased from a reputed vendor of herbal Medical products, Everest Marunthakam, Usan, Mirusuvil (Registered at Sri Lanka Ayurveda Medical Council). marketed formuation of Thalangai ennai (MFT) was standardized based on their organoleptic characters (physical characteristic), physicochemical properties (pH. specific gravity) and thin chromatography (TLC) according to standard procedures and compared with the Thalangai ennai of Drug Unit of Indigenous Medicine, Jaffna (DUT).

Ingredients and preparation method of Thalangai ennai

Botanically pure and authentic ingredients were used in the preparation of *Thalangai ennai* accordance with the guidelines stated in the Siddha Literature.

Organoleptic evaluation

Both samples of oil (Thalangai ennai) and its water and dichloromethane extracts were subjected to the organoleptic characterization such as appearance, touch, colour, clarity and odour and compared with the values of Sesame oil (Which is used as a base in preparing these oils) separately. The organoleptic characters of the samples were evaluated based on the method described by Siddiqui et al. 1995.

Physico-chemical investigations

Both samples of oil (*Thalangai ennai*) were subjected for determination of preliminary physicochemical parameters such as pH value and specific gravity.

Development of Thin-layer chromatography (TLC) fingerprints of Thalangai ennai

Thin Layer Chromatography (TLC) finger print of the *Thalangai ennai* was studied after dissolving the oil in water and dichloromethane.

Five µl of dichloromethane extract of each oil (MFT & DUT) was spotted on to TLC plates (8.5 x 5.3 cm) coated with silica gel (pre-coated, GF25a) and separated using a variety of solvent system. Different solvent systems were used for separation. Although the separation of the extracts occurred in the solvent system comprised of hexane: dichloromethane: 1% methanol in dichloromethane (1:4:5 v/v) as the mobile phase, the best separation of the extracts occurred in the solvent system comprised of pure dichloromethane as the mobile phase. After development visible spots were not found for each oil extract. Numbers of spots were observed under day light and UV light (254 & 366 nm). Visualization was

attempted by spraying with vanillin sulphate reagent for each oil extract and heating the plate for 5-10 minute (100-105°C). The colour and Rf values

Rf=

of the spots were recorded carefully and the chromatogram was documented by graphical copying (WHO, 1998).

Distance travelled by solute

Distance travelled by solvent

Results and Discussion

standards physicochemical The available for the standardization of Siddha Medicated oils are insufficient. The analytical values available in the pharmacopoeia standards for siddha formulations are not finger print standards for each of oil. The first step current the towards this goal. preliminary investigation was generate data on undertaken to physico-chemical properties, including and characters. organoleptic chromatographic profiles to determine the quality and purity of Thalangai ennai. It is very difficult to perform a study on Siddha medicated oils which has a large number of herbs used in the formulation. Hence in this study only Thalangai ennai was selected. As there is not any evidence for detailed physicochemical and TLC evaluation on *Thalangai ennai* is reported. Therefore present work is taken up in the view to standardize the *Thalangai ennai*. In the present study different brands of *Thalangai ennai* was subjected to physicochemical parameters, organoleptic characters and TLC finger printing for preliminary standardization.

Organoleptic characters and physicochemical properties

Table 1, summarizes the organoleptic properties, pH values and Specific gravity of the Marketed formulation (MFT) and Drug Unit of Indigenous Medicine (DUT) *Thalangai Ennai* with Sesame oil at room temperature and Table 2 summarizes the organoleptic properties of its water and dichloromethane

Table 1: Analytical parameters of Thalangai Ennai MFT & DUT with Sesame oil

Parameters	Thalangai ennai (MFT)	Thalangai ennai (DUT)	Sesame Oil
Appearance	Viscous	Viscous	Viscous
Touch	Oily	Oily	Oily
Colour	Red	Golden brown	Brown
Clarity	Clear	Clear	Clear
Odour	Characteristic	Characteristic	Characteristic
pH value (29°C)	4.97	4.61	4.89
Specific gravity	0.9173	0.9185	0.923

As seen in Table 1, both Brands of the *Thalangai ennai* had similar organoleptic properties except for the colour of the each oil compared with Sesame oil. The results showed that, the specific gravity and pH value for *Thalangai ennai* DUT were found to be

0.9185 and 4.61; in case of MFT these were found to be 0.9173 and 4.97 at room temperature (29°C) respectively. The pH and specific gravity of these brands were showed not remarkable difference between them.

Table 2: Organoleptic properties of the water and dichloromethane extracts of the of *Thalangai ennai* MFT & DUT

Parameter	MFT		DUT	
\$	Water extract	Dichlorometha ne extract	Water extract	Dichlorometha ne extract
Appearance	Liquid	Liquid	Liquid	Liquid
Touch	Liquid	Liquid	Liquid	Liquid
Colour	Colourless	Colourless	Pale yellow	Light yellow
Clarity Odour	Clear Characteristic	Clear Characteristic	Clear Characteristic	Clear Characteristic

As seen in Table 2, both extracts of the each *Thalangai ennai* had similar organoleptic properties except for the colour of the water and dichloromethane extracts of oils.

Thin Layer Chromatography (TLC) finger print analysis

The numbers of different solvent systems were employed to generate fingerprint profiles for these oils. The TLC finger print of both the formulations were comparable and revealed dichloromethane extract of the oil samples showed twelve spots with Rf values ranging from 0.02 to 0.97 in the hexane: dichloromethane: 1% methanol in dichloromethane (1: 4: 5 v/v) solvent system for both brands and fifteen spots with Rf values ranging

from 0.03 to 0.96 (MFT) and 0.03 to 0.97 (DUT) in the pure dichloromethane solvent system.

Although the separation of the oil extracts occurred in the solvent system comprised of hexane: dichloromethane: 1% methanol in dichloromethane (1:4:5 v/ v) as the mobile phase, the best separation of the oil extracts occurred in the solvent system comprised of pure dichloromethane (v) the mobile phase. dichloromethane extract of different brands of Thalangai ennai (MFT and DUT) analyzed by TLC, the number of spots with the Rf values were observed not remarkable differences in between these both oil.




Figure 1: TLC fingerprints of oil extract a) 254 nm, b) 366 nm and c) visible after spray in the Solvent system of Hexane: Dichloromethane: 1% Methanol in Dichloromethane (1: 4: 5 v/v) mixture




Figure 2: TLC fingerprints of oil extract a) 254 nm, b) 366 nm and c) visible after spray in the Solvent system pure Dichloromethane

Conclusion

The results obtained with the both brands of *Thalangai ennai* (MFT and DUT) were found to be comparable and variations were not remarkable. These set of parameters presented in this paper can be used as reference standards for the quality assurance of *Thalangai ennai*. Though the analytical values can be as preliminary reference

standards, these values mostly related to the purity of the sesame oil, the qualitative and quantitative estimation of each single ingredient of *Thalangai* ennai requires further research work in future.

Acknowledgement: The financial support provided by the Higher Education Twenty-first Century

(HETC) Project, Ministry of Higher Education, Nugegoda, Sri Lanka (Reference No - JFN/ Sidda/ N1), and the guidance given by Dr (Mrs.) Menuka Arawwawala, Industrial Technology Institute (ITI), Colombo 07, to conduct the physicochemical and TLC finger print studies are gratefully acknowledged. This work is a part of the research carried for postgraduate programme (PhD) at faculty of Graduate Studies, University of Jaffna.

References

Hepsibah PTA, Prasad NBR & Sanjeev Kumar P, 1998. "Standardization of Ayurvedic oils". Ancient Science of Life; 17(4): pp. 280 – 283.

Kumar A and Baxi AJ, 2007. "Standardization of Ayurvedic Medicated Oil and the effect of Moorchhan on the amount of Marker in the oil", Indian Drugs, 44(2), pp. 123-127.

Organization Mondiale De La Sante, 1998. Quality control methods for plant materials, 559, Rev.1, Original English, World Health Organization (W.H.O.), P. 8 - 67.

Ponniahpillai, I., 2000. "Vaithiya vihazham ennum Amirthasakara pathartha sudamanium vaidhya thelivum (Irupalai shettiyar)", Provincial Department of Indigenous Medicine, Ministry of Health, N.E.P., 35-36.

Siddiqui, Hakim MA. 1995. "Format for the pharmacopoeial analytical standards of compound formulation, workshop on standardization of Unani drugs, (appendix), 24-25 January, New Delhi: Central Council for Research in Unani Medicine (CCRUM).

Thirunarayanan. T, 1988. "An Introduction to Siddha Medicine", Thirukumaran Publishers, Tiruchendur, 7-14.