

Growth and Yield response of Maize (*Zea mays*) intercropped with different crop densities of Green gram (*Vigna radiate*)

Kirija,M., Sivachandiran,S., Pradheeban,L., Selvanathan,N.

gisajdfjgjasjdgfasdgfoiuj

Abstract

Impact of weed control methods on the persistence of *Partheniumhysterophorus* (L), and quantity and quality of tomato (*Solanum lycopersicum* L) yield was studied in two farmer fields i.e. a site heavily infested with *P. hysterophorus* (Parthenium-site) and Parthenium-free site, in the Jaffna peninsula of the Northern Province of Sri Lanka. The split-plot design consisted of two main plots i.e. manual weeding and ploughing (both followed by harrowing), and three sub-plots i.e. use of a pre-emergent herbicide (Oxyfluorfen; 240 g/L EC at 2 days before planting), mulching [*Gliricidiasepium* (Jacq) Kunth ex Walp. leaves at 12 kg per plot on fresh weight basis], and un-weeded plot after crop establishment (control), in three replicates. The soil seed bank of the Parthenium-site was dominated by *P. hysterophorus* as expected, and the Parthenium-free site was dominated by the perennial sedge *Cyperusrotundus* (L). At tomato harvest, the un-weeded sub-plots recorded the highest weed densities ($p<0.05$) when compared to the rest, The Parthenium-site showed a higher weed dry weight ($p<0.05$) when compared to that of the Parthenium-free site, due to higher emergence of the Parthenium weed in the former. Mulching was the best sub-plot treatment for Parthenium control in combination with the main plot treatments. The average fruit weight of tomato at the Parthenium-free site was 7.8% higher than that of the Parthenium-site (47.8 g per fruit). In the Parthenium-site, mulching resulted in a 6.4% higher fruit weight and 58% higher total yield ($p<0.05$) when compared to rest of the sub-plot treatments. The weed competition negatively affected the tomato yield ($Y=-0.7551X+7.88$; $R^2=0.58$; $p<0.05$), with Parthenium weed playing a dominant role. Mulching with *G. sepium* coupled with manual weeding during land preparation or ploughing would suppress growth and development of weeds including *P. hysterophorus* and enhance yield of tomato.